

TITLE: SINGLE KINK DYNAMICS IN AN EASY-PLANE CLASSICAL ANTIFERROMAGNET  
CHAIN

AUTHOR(S): Gary M. Wysin  
Alan R. Bishop  
Jaan Oitmaa

SUBMITTED TO: Journal of Physics C.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy

---

Los Alamos Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

SINGLE KINK DYNAMICS IN AN EASY-PLANE

CLASSICAL ANTIFERROMAGNET CHAIN

G. M. Wysin<sup>†</sup> and A. R. Bishop  
Center for Nonlinear Studies and Theoretical Division  
Los Alamos National Laboratory  
Los Alamos, NM 87545, U.S.A.

and

J. Oitmaa  
School of Physics, The University of New South Wales  
P.O. Box 1, Kensington, NSW 2033  
AUSTRALIA

J. Phys. C 19, 221 (1986).

<sup>†</sup>Permanent address: LASSP, Cornell University, Ithaca, NY 14853.

## Abstract

We present an Ansatz for kink excitations in an easy-plane classical antiferromagnetic chain with a magnetic field in the easy plane. The utility of the Ansatz is that it presents the in-plane (XY) and out-of-plane (YZ) kinks as belonging to one continuously connected energy dispersion curve. Linear stability analysis applied to YZ kinks shows that there is a velocity dependent critical field necessary for stability. We also present results of a numerical integration of the equations of motion which verifies the YZ kink stability regimes, as well as showing that XY kinks are stable over a wide range of fields and velocities.

## I. Introduction

Recently there has been considerable experimental and theoretical interest in the low temperature properties of 1-D easy-plane anti-ferromagnets in the presence of magnetic fields applied in the easy plane. A typical Hamiltonian assumed to describe these materials is (Mikeska 1980):

$$H = \sum_n \{ J \vec{S}_n \cdot \vec{S}_{n+1} + A(S_n^z)^2 - g\mu_B B_x S_n^x \} , \quad (1)$$

where the notation is standard and  $J > 0$ ;  $A > 0$  represents the effect of dipole-dipole interactions, and the field  $B_x$  is in the easy (XY) plane. Neutron scattering experiments (Heilmann et al. 1979, Boucher et al. 1982) on the  $S = 5/2$  compound  $(CD_3)_4 NMn Cl_3$  (TMMC), an example of this type of material, have shown an interesting behavior of the in-plane and out-of-plane spin wave dispersions measured as functions of the applied field. In particular, there is evidence for a crossover field, where these spin wave energies become equal, which can be considered to be a switching of the hard anisotropy axis from the dipole anisotropy axis to an axis parallel to the field (Harada et al. 1981).

While this hard axis switching arises simply in a linear spin wave theory, it is also manifested in the nonlinear regime (Harada et al. 1981, Flüggen and Mikeska 1983). The nonlinear dynamics in the continuum limit has been mapped approximately to the sine-Gordon equation, with kink or "soliton" excitations being  $\pi$  rotations of the spins either in the easy plane (XY kinks) or in a plane whose normal is parallel to the field (YZ kinks). These have static energies  $g\mu_B B_x S$  and  $(8AJS^4)^{1/2}$ , so that the critical field at which the two energies are equal is  $B_c = (8AJS^2)^{1/2}/(g\mu_B)$ . This is the same field for which the in-plane and out-of-plane spin waves (at the zone boundary) have equal energies.

The XY and YZ kinks have previously been considered as distinct special solutions of the continuum equations of motion resulting from (1), belonging to two distinct branches in energy dispersion. Through the application of a variational Ansatz (Wysin 1985), we show here that they are actually just two limits of a single continuously connected dispersion curve. The Ansatz (below) is motivated by a corresponding classical ferromagnet Ansatz used by Liebman et al. (1983). The spins are assumed to rotate in a plane tilted at some angle to the XY plane, with a sine-Gordon motion of the spins in that plane. General features of results derived from this Ansatz are very similar to those for the ferromagnet -- including the existence of a maximum XY kink velocity which is less than the spin wave velocity (see also Wysin et al. 1982, 1984). There is an important difference, however, between the ferromagnetic and antiferromagnetic Ansatz: For the ferromagnet, the ground state is unique,  $\vec{S} = (1,0,0)$ , while for the antiferromagnet, the ground state is two fold degenerate, with  $\vec{S}_{\text{odd}} = (\frac{1}{2}\beta, \sqrt{1-\frac{1}{16}\beta^2}, 0)$ ,  $\vec{S}_{\text{even}} = (\frac{1}{2}\beta, -\sqrt{1-\frac{1}{16}\beta^2}, 0)$  or vice versa, with  $\beta \equiv g\mu_B B_x / JS$ . In any antiferromagnet Ansatz, the spin at a given lattice site must rotate from one ground state to the other. The fact that the two ground state sublattices are not exactly antiparallel (due to the spin flop effect) leads to considerable complications which cannot be ignored. We should note that for an easy-plane ferromagnet with an Ising symmetry-breaking field ( $J < 0$  and  $\beta_x S_n^x$  replaced by  $\beta_x (S_n^x)^2$  in Eq. (1)) the two degenerate ground states are exactly antiparallel, and this Ansatz precisely reproduces the soliton solutions given by Sklyanin (1979).

We have also considered the stability of kink structures in these systems. This has been approached in two ways: (i) The stability can

be tested by numerical integration of the discrete equations of motion. We have used an initial kink profile obtained from either the XY or YZ sine-Gordon limits of the equations of motion, and followed the time evolution. Alternatively, one can use an initial kink profile as obtained from the Ansatz presented here. Details are given below; or (ii) A linearized stability analysis about sine-Gordon static kink profiles can be used. This can be followed through analytically for the YZ static kinks, but we have not been able to generally solve the stability equations for the XY static kink. We find from both approaches that static YZ kinks are stable only if  $B > B_c$ ; there is no corresponding instability field for static XY kinks -- they are always stable over a wide range of applied field. For moving YZ kinks one finds from the numerical simulations that the stability can either be enhanced or diminished by the motion, depending on whether the motion causes the spins to cant further toward the field or away from it. The linear stability analysis confirms this behavior. The numerical simulations indicate that dynamic XY kinks are always stable.

## II. The continuum sine-Gordon limits

First we consider the continuum equations of motion resulting from (1), and the XY and YZ sine-Gordon limits, as given by Flüggen and Mikeska (1983). The spins at the even or odd lattice sites, referred to as the A and B sublattices, have been parametrized in terms of four angles in spherical polar coordinates:

$$\vec{S}_{A/B} = \pm S(\sin(\theta \pm \theta) \cos(\phi \pm \phi), \sin(\theta \pm \theta) \sin(\phi \pm \phi), \cos(\theta \pm \theta)) . \quad (2)$$

S is the length of the spin vectors, and  $\theta$ ,  $\theta$ ,  $\phi$  and  $\phi$  are all functions

of time and position  $x$  on the chain. In order to take the continuum limit, we must assume  $\theta, \phi \ll 1$ , as well as slow spatial variations in  $\phi$  and  $\theta$ . Measuring time in units of  $\hbar/JS$ , and using the lattice constant as the length unit, the evolution equations are:

$$(\phi_{xx} - \frac{1}{2}\ddot{\phi})\sin\theta + 2(\theta_x \phi_x - \frac{1}{2}\dot{\theta}\dot{\phi})\cos\theta = -\frac{1}{2}\beta^2 \sin\theta \sin\phi \cos\phi + \frac{1}{2}\beta\dot{\theta} \sin\theta \cos\phi \quad (3a)$$

$$(\theta_{xx} - \frac{1}{2}\ddot{\theta})\csc\theta - (\phi_x^2 - \frac{1}{2}\dot{\phi}^2)\cos\theta = (\frac{1}{2}\beta^2 \cos^2\phi - \alpha)\cos\theta - \frac{1}{2}\beta\dot{\phi}\sin\theta \cos\phi, \quad (3b)$$

where  $\beta \equiv g\mu_B B_x/JS$  and  $\alpha \equiv 2A/J$ , and space and time derivatives are denoted by subscripts and dots, respectively. The "small" angles  $\theta$  and  $\phi$  are given by:

$$\theta = \frac{1}{2}(\beta \cos\theta \cos\phi - \dot{\phi} \sin\theta) \quad (4a)$$

$$\phi = \frac{1}{2} \csc\theta(\dot{\theta} - \beta \sin\phi). \quad (4b)$$

Here we have assumed  $\alpha \ll 4$ , which is the case for most real materials ( $\alpha \cong .04$  for TMMC).

#### IIA. XY kinks

Statically, Eq. (3b) is satisfied exactly when  $\theta = \frac{1}{2}\pi$ . Equation (3a) then becomes a static sine-Gordon equation for the variable  $\psi \equiv 2\phi - \pi$ . More generally, one can assume  $\theta = \frac{1}{2}\pi - \theta_s$ , where  $\theta_s \ll 1$ , and linearize the equations in  $\theta_s$ : this yields a dynamic sine-Gordon equation for  $\psi$ :

$$\psi_{xx} - \frac{1}{2}\ddot{\psi} = \frac{1}{2}\beta^2 \sin\psi \quad (5a)$$

and

$$(\theta_{sxx} - \frac{1}{2}\ddot{\theta}_s) + (\phi_x^2 - \frac{1}{2}\dot{\phi}^2 + \frac{1}{2}\beta^2 \cos^2\phi - \alpha)\theta_s = \frac{1}{2}\beta\dot{\phi} \cos\phi. \quad (5b)$$

A kink solution to Eqs. (5), for  $\beta^2 \ll \alpha$ , is a  $\pi$  rotation in the XY plane, moving as a traveling-wave at velocity  $v$ :

$$\Phi = \frac{\pi}{2} + 2 \tan^{-1} \exp(\tilde{z}); \quad \tilde{z} \equiv \frac{1}{2}\beta\gamma(x-vt) \quad (6a)$$

$$\theta_s = -\frac{1}{2}(\beta^2\gamma v/\alpha) \operatorname{sech}^2(\tilde{z}); \quad \gamma \equiv (1-\frac{1}{2}v^2)^{-\frac{1}{2}}. \quad (6b)$$

Note that the spin wave velocity  $c = 2$  in these units. Although solution (6) is valid only at low fields, we have used these profiles as initial conditions in numerical simulations for arbitrary  $\beta$ . The energy of the kink in Eq. (6) is  $E_{XY} = \beta\gamma JS^2$  plus corrections which depend on  $\beta^2/\alpha$ . There is considerable experimental evidence in TMMC demonstrating the existence of these modes (Boucher and Renard 1980, Boucher et al. 1983, 1984, Regnault et al. 1982).

## II.B. YZ kinks

Equation (3a) is satisfied exactly by taking  $\Phi = \frac{1}{2}\pi$ , in which case (3b) becomes a sine-Gordon equation in the variable  $2\theta - \pi$ . The dynamic kink solution is now a  $\pi$  rotation in the YZ plane:

$$\theta = \frac{\pi}{2} + 2 \tan^{-1} \exp(z); \quad z \equiv \gamma\sqrt{\alpha}(x-vt). \quad (7)$$

Although result (7) has been obtained "exactly" within the small angle approximations  $\theta, \phi \ll 1$ , substituting (7) into (4) one finds  $\theta = 0$ , but

$$\phi = \frac{1}{2}(\gamma\sqrt{\alpha}v \operatorname{sech}z + \beta)\coth z. \quad (8)$$

Thus at the center of the kink ( $z=0$ ),  $\phi$  is no longer small as has been assumed, and this solution is not valid. The profile has an unphysical cusp in  $S^x$  which prohibits its use as initial conditions in numerical simulations.



One can correct this problem, and still obtain an exact sine-Gordon equation for YZ kinks, by using a more appropriate coordinate system. Spherical polar coordinates, where the x-axis (i.e., the field direction) is the polar axis, exploit the symmetry of the YZ kinks. With this motivation, we have re-parametrized the spins in terms of new angles  $\theta$ ,  $\theta$ ,  $\phi$ ,  $\phi$  in x-polar spherical coordinates:

$$\begin{matrix} \vec{S} \\ A \\ B \end{matrix} = \pm S(\cos(\theta \pm \theta), \sin(\theta \pm \theta) \cos(\phi \pm \phi), \sin(\theta \pm \theta) \sin(\phi \pm \phi)) . \quad (9)$$

The dynamical equations for the four variables now become (Wysin 1985)

$$\begin{aligned} \dot{\theta} &= 4\phi \sin\theta + \alpha(\phi \sin\theta \cos 2\phi + \frac{1}{2}\theta \cos\theta \sin 2\phi) \\ \dot{\phi} &= -4\theta \csc\theta + \alpha(\theta \sin\theta \sin^2\phi - \phi \cos\theta \sin 2\phi) - \beta \\ \ddot{\theta} &= -(2\phi_{xx} \theta + 4\phi\theta) \cos\theta - \phi_{xx} \sin\theta + \frac{1}{2}\alpha \sin\theta \sin 2\phi \\ \ddot{\phi} &= -(4\phi^2 + \phi_x^2 - 4\theta^2 \csc^2\theta) \cos\theta + \theta_{xx} \csc\theta - \alpha \cos\theta \sin^2\phi . \end{aligned} \quad (11)$$

Elimination of the small angles  $\theta$ ,  $\phi \ll 1$ , under the assumption  $\alpha \ll 4$ , leads to equations similar in structure to Eqs. (3) and (4):

$$(\phi_{xx} - \frac{1}{2}\ddot{\phi}) \sin\theta + 2(\theta_{xx} \phi - \frac{1}{2}\dot{\theta}\dot{\phi}) \cos\theta = \frac{1}{2}\alpha \sin\theta \sin 2\phi + \frac{1}{2}\beta \dot{\theta} \cos\theta \quad (12a)$$

$$(\theta_{xx} - \frac{1}{2}\ddot{\theta}) \csc\theta - (\phi_x^2 - \frac{1}{2}\dot{\phi}^2) \cos\theta = (\alpha \sin^2\phi - \frac{1}{2}\beta^2) \cos\theta - \frac{1}{2}\beta \dot{\phi} \cos\theta \quad (12b)$$

$$\phi = \frac{1}{2}\dot{\theta} \csc\theta \quad (13a)$$

$$\theta = -\frac{1}{2}(\beta + \dot{\phi}) \sin\theta . \quad (13b)$$

The YZ kink is now found by taking  $\theta = \frac{1}{2}\pi$ , which satisfies (12b) exactly, and then (12a) becomes a sine-Gordon equation for  $2\phi$ , with kink solution

$$\phi = 2 \tan^{-1} \exp z; \quad \phi = 0 \quad (14a)$$

$$\theta = -\frac{1}{2}(\beta - \gamma\sqrt{\alpha}v \operatorname{sech} z) . \quad (14b)$$

We now have smooth functions for  $\theta$  and  $\phi$ , and we have used this profile as initial conditions for numerical simulations. The kink in Eq. (14) has energy

$$E_{YZ} = 2\gamma\sqrt{\alpha}(1 - \frac{\beta^2}{32}(1 - \frac{1}{2}v^2)).$$

### IIC. YZ stability analysis

The stability of the YZ kink solution to Eq. (12) can be determined by assuming small perturbations  $\tilde{\theta}, \tilde{\theta}, \tilde{\phi}, \tilde{\phi}$  about the sine-Gordon solution  $\theta_0, \theta_0, \phi_0, \phi_0$ , substituting into (11) and linearizing in these perturbations. The small angles  $\tilde{\theta}, \tilde{\phi}$  can be eliminated from the resulting equations, and we obtain decoupled equations for  $\tilde{\theta}, \tilde{\phi}$ . For  $\alpha \ll 4$ , they are

$$\tilde{\theta}_{xx} + (\phi_{0x}^2 - 4\theta_0^2 + \alpha \sin^2 \phi_0) \tilde{\theta} = \frac{1}{2} \ddot{\tilde{\theta}} \quad (15a)$$

$$\tilde{\phi}_{xx} - \alpha \cos 2\phi_0 \tilde{\phi} = \frac{1}{2} \ddot{\tilde{\phi}} . \quad (15b)$$

We assume  $\tilde{\theta} \sim e^{i\omega_1 t}$ ,  $\tilde{\phi} \sim e^{i\omega_2 t}$  time dependences, and using Eqs. (14), (15) yields the eigenvalue problems (Wysin 1985)

$$-\gamma^2 \tilde{\theta}_{zz} + (1-2 \operatorname{sech}^2 z - \frac{\beta\gamma v}{2\sqrt{\alpha}} \operatorname{sech} z) \tilde{\theta} = \lambda_1 \tilde{\theta} \quad (16a)$$

$$-\gamma^2 \tilde{\phi}_{zz} + (1-2 \operatorname{sech}^2 z) \tilde{\phi} = \lambda_2 \tilde{\phi} \quad (16b)$$

with

$$\omega_1^2 = 4\alpha(\lambda_1 - 1) + \beta^2 \quad (16c)$$

$$\omega_2^2 = 4\alpha\lambda_2 \quad (16d)$$

For a zero velocity kink, both equations have  $\operatorname{sech}(z)$  bound state solutions, with eigenvalues  $\lambda_1 = \lambda_2 = 0$ . The corresponding eigenfrequencies are

$$\omega_1^2 = \beta^2 - 4\alpha \quad (17)$$

$$\omega_2^2 = 0 .$$

The bound state associated with the  $\tilde{\phi}$  variable is the Goldstone translation mode and involves no structural instability. The bound state for  $\tilde{\theta}$ , however, can have an imaginary eigenfrequency indicating an instability.<sup>++</sup> Specifically, for fields  $\beta^2 < 4\alpha$ , static YZ kinks are unstable, and the mode responsible for this instability causes the spins to tilt away from the z-axis. This result has been confirmed by numerical integration of the discrete equations of motion. Starting from a static YZ kink profile, the spins move toward a configuration involving a lower energy XY kink, with the excess energy given to spin waves. Conversely, for  $\beta^2 > 4\alpha$ , static YZ kinks have been found numerically to be stable.

---

<sup>++</sup>The analysis by Flüggen and Mikska (1983) did not reveal this instability due to their use of z-polar spherical coordinates.

For nonzero  $v$ , the potential for  $\tilde{\theta}$  is modified, and therefore so is the bound state eigenvalue  $\lambda_1$ . If  $v$  is positive, then the potential becomes deeper, and therefore  $\lambda_1$  becomes less than zero. From (16c), the field necessary for the kink to be stable is now  $\beta^2 > 4\alpha(1-\lambda_1)$ ; that is, a kink with positive velocity requires a larger field to be stable than does the static kink. For a kink with negative velocity, the opposite is true -- a smaller field is necessary. Even if  $\beta^2 < 4\alpha$ , there can be stable moving YZ kinks. This has been verified by numerical simulation.

These comments can be made quantitative by estimating  $\lambda_1(v)$  using perturbation theory. A simple first order calculation gives  $\lambda_1(v) \cong -(\pi\beta/4\alpha^{3/2})(v/c)$ ; to get a result correct to order  $(v/c)^2$  one needs to use second order perturbation theory (Wysin 1985). To first order in  $v/c$ , then, the stability criterion  $\omega_1^2 > 0$  for moving YZ kinks becomes  $v/c < (\beta^2 - 4\alpha)/(\pi\beta\alpha^{3/2})$ . The numerical simulation (section IV) gives results consistent with this relationship.

It should be noted that this direction dependent stability does not really imply a preferred direction of motion for YZ solitons, since we have presented the stability arguments only for one of the four possible soliton solutions (there are two kinks and two antikinks possible). The conclusion that can be drawn, however, is that those YZ solitons in which the motion causes the spins to cant further toward the field (compared to the ground state canting) have increased stability.

### III. The Variational Ansatz

In the above discussion the XY and YZ kinks have been assumed to be two independent types of excitations. For a dynamic XY kink, the spins of the A sublattice rotate approximately in a plane tilted at an

angle  $\theta_A$  to the XY plane, while the spins of the B sublattice move in a similar plane which may be at a slightly different angle  $\theta_B$ . For low velocity kinks  $\theta_A$  and  $\theta_B$  are near zero, and as the velocity increases so do  $\theta_A$  and  $\theta_B$ . For dynamic YZ kinks,  $\theta_A$  and  $\theta_B$  are near  $\frac{1}{2}\pi$ , and should approximately satisfy  $\theta_A = \frac{1}{2}\pi - \delta$ ,  $\theta_B = \frac{1}{2}\pi + \delta$ , where  $\delta$  is a small parameter. (In fact, the Ansatz predicts that  $\delta$  is proportional to the velocity.) These observations suggest that one can make an Ansatz for a general kink excitation which will include both the XY and YZ kinks as special limiting cases. The Ansatz will involve variational parameters  $\theta_A$ ,  $\theta_B$  and a width  $w$ , these parameters being determined by extremizing the Lagrangian with respect to the parameters. This Ansatz calculation closely parallels a similar one for the corresponding ferromagnet (Liebmann et al 1983).

An appropriate Ansatz for the xyz components of unit length spins,  $\hat{o}_A$ ,  $\hat{o}_B$ , is constructed as follows: Consider the spins on the A sublattice first. The trajectory on the unit sphere must be an arc starting from one ground state  $(\frac{1}{2}\beta, -(1-\beta^2/16)^{\frac{1}{2}}, 0)$ , passing through a point  $(\cos q, 0, \sin q)$  at the center of the kink, and then ending at the opposite ground state  $(\frac{1}{2}\beta, +(1-\beta^2/16)^{\frac{1}{2}}, 0)$ . The plane in which the tips of the spin vectors move is tilted on an angle  $\theta_A$ , such that  $\tan\theta_A = \sin q / (\cos q - \frac{1}{2}\beta)$  (see Fig. 1). In the  $x'y'z'$  coordinate system, obtained by rotating the xyz system through  $\theta_A$  about the y axis, the equation of this plane is simply  $z' = -\frac{1}{2}\beta \sin\theta_A$ . In order to determine the  $x'$  and  $y'$  components of this trajectory, consider the sum and difference vectors  $\vec{M} = \frac{1}{2}(\hat{o}_A + \hat{o}_B)$ ,  $\vec{N} = \frac{1}{2}(\hat{o}_A - \hat{o}_B)$  (supposing for the moment that the B sublattice moves in the same plane). The vector  $\vec{M}$  approximately traces out a circle of radius  $r_M = \frac{1}{8}\beta \cos\theta_A$ , with center at

$(x', y') = (\frac{1}{8} \beta \cos \theta_A, 0)$ . The vector  $\vec{N}$  approximately traces out a semi-circle of radius  $r_N = (1 - \beta^2/16)^{1/2}$ , centered at  $(x', y') = (0, 0)$ . Therefore we take (see Fig. 2)

$$M_{x'} = r_M (1 + \cos \phi_{sG}) = 2r_M \tanh^2 z \quad (18a)$$

$$M_{y'} = r_M \sin \phi_{sG} = -2r_M \tanh z \operatorname{sech} z \quad (18b)$$

$$N_{x'} = r_N \sin(\frac{1}{2} \phi_{sG}) = r_N \operatorname{sech} z \quad (18c)$$

$$N_{y'} = -r_N \cos(\frac{1}{2} \phi_{sG}) = r_N \tanh z \quad (18d)$$

where

$$\phi_{sG} = 4 \tan^{-1} \exp(z) ; z = \frac{x-vt}{w} \quad (19)$$

Transforming back to the  $\hat{o}$  variables, it is necessary to multiply the  $x'y'$  components by a factor  $f_A(z) = (1 - \sin^2 \epsilon_A \operatorname{sech}^2 z)^{-1/2}$ , in order to keep the spin length fixed ( $\epsilon_A$  is shown in Fig. 1). If we assume now that the A and B sublattices rotate in planes at different angles  $\theta_A$  and  $\theta_B$ , then we can write the Ansatz as follows:

$$\begin{aligned} \sigma_{Ax'} &= f_A(z) (M_{x'} + N_{x'}) , & \sigma_{Bx''} &= f_B(z) (M_{x''} - N_{x''}) , \\ \sigma_{Ay'} &= f_A(z) (M_{y'} + N_{y'}) , & \sigma_{By''} &= f_B(z) (M_{y''} - N_{y''}) , \\ \sigma_{Az'} &= -\frac{1}{2} \beta \sin \theta_A , & \sigma_{Bz''} &= -\frac{1}{2} \beta \sin \theta_B . \end{aligned} \quad (20)$$

Here single primes refer to the "A" coordinate system at angle  $\theta_A$ , while double primes refer to the "B" coordinate system at angle  $\theta_B$ . The single function  $\phi_{sG}$  describes the distribution of the spins on the trajectories for both sublattices.

Using energy units  $JS^2$ , and time units  $\hbar/JS$ , the continuum limit Lagrangian is:

$$L = \int dx \left\{ \frac{1}{2} \sum_{i=A,B} \left[ \sigma_i^z \frac{d}{dt} \tan^{-1} \left( \frac{\sigma_i^y}{\sigma_i^x} \right) - \frac{1}{2} \alpha (\sigma_i^z)^2 + \frac{1}{2} \beta \sigma_i^x \right] - (\hat{\sigma}_A \cdot \hat{\sigma}_B - \frac{1}{2} \frac{d\hat{\sigma}_A}{dx} \cdot \frac{d\hat{\sigma}_B}{dx}) \right\} . \quad (21)$$

The first term in Eq. (21) is simplified by changing  $\sigma_A^z \rightarrow \sigma_A^z - 1$  and  $\sigma_B^z \rightarrow \sigma_B^z + 1$ , which changes the Lagrangian by an additive constant that depends only on  $\beta$ , and removes unphysical step functions at  $\theta_A = \frac{1}{2}\pi$  and  $\theta_B = -\frac{1}{2}\pi$ . Since we minimize  $L$  with respect to the variational parameters  $w$ ,  $\theta_A$  and  $\theta_B$ , this constant does not affect any of the results that follow. After rotating the Ansatz back to the original xyz coordinate system, and considering  $\sqrt{\alpha}$ ,  $\beta$  and  $\Delta \equiv \theta_B - \theta_A$  as small parameters, the Lagrangian (relative to the ground state) is found to be (Wysin 1985):

$$L(\theta_A, \Delta, w) = vK(\theta_A, \Delta) - \frac{1}{w} F(\theta_A, \Delta) - wG(\theta_A, \Delta) , \quad (22a)$$

where

$$F(\theta_A, \Delta) = (r_N^2 - \frac{\beta^2}{48} \cos^2 \theta_A) - (\frac{\pi\beta}{16} r_N \sin \theta_A) \Delta - \frac{1}{6} \Delta^2 \quad (22b)$$

$$G(\theta_A, \Delta) = (\alpha r_N^2 \sin^2 \theta_A + \frac{1}{2} \beta^2 \cos^2 \theta_A) + \frac{1}{2} (\alpha r_N^2 - \frac{1}{2} \beta^2) \sin 2\theta_A \cdot \Delta + \Delta^2 \quad (22c)$$

$$K(\theta_A, \Delta) = \pi + \Delta + \frac{\pi\beta}{4} (\sin \theta_A + \frac{1}{2} \Delta \cos \theta_A) \equiv k_0 + k_1 \Delta . \quad (22d)$$

Minimization of  $L$  with respect to the width  $w$  leads to

